

ABCD Human Subjects Study

Adolescent Brain Cognitive Development – ABCDSTUDY.org

Release Notes: Adolescent Brain Cognitive Development StudySM (ABCD) Data Release 3.0

Resting-State Functional Magnetic Resonance Imaging (rs-fMRI)

<http://dx.doi.org/10.15154/1519007>

October 2020

Change Log

October 2020 – Data Release 3.0

- Initial release

List of Instruments

Name of Instrument	Short Name
ABCD rsfMRI Gordon Network Correlations	abcd_betnet02
ABCD rsfMRI Network to Subcortical ROI Correlations	mrirscor02
ABCD rsfMRI Temporal Variance	abcd_mrirstv02
ABCD rsfMRI Destrieux	abcd_mrirsfd01

General Information

The following information refers to the Adolescent Brain Cognitive DevelopmentSM (ABCD) Study Data Release 3.0 available from <https://nda.nih.gov/abcd>. An overview of the ABCD Study[®] is at <https://abcdstudy.org> and detailed descriptions of elements of the study are listed in <https://abcdstudy.org/scientists/protocols>.

This document describes the contents of various instruments available for download. To understand the context of this information, see *Release Notes ABCD README FIRST* and *Release Notes ABCD Imaging Instruments*.

- Image types
 - field maps: spin echo images with opposite phase encode polarity
 - multi-frame, gradient echo, echo-planar imaging
- Image processing (common to all fMRI)
 - head motion corrected by registering each frame to the first using AFNI's 3dvolreg (Cox, 1996)
 - B₀ distortions were corrected using the reversing gradient method with FSL's TOPUP (Andersson et al., 2003, Smith et al., 2004)
 - displacement field estimated from spin-echo field map scans
 - applied to gradient-echo images after adjustment for between-scan head motion
 - corrected for gradient nonlinearity distortions (Jovicich, et al., 2006)
 - between scan motion correction across all fMRI scans in imaging event
 - registration between T₂-weighted, spin-echo B₀ calibration scans and T₁-weighted structural images performed using mutual information (Wells, et al., 1996)
- rs-fMRI specific pre-processing -- NOTE: not included in "minimal processing"
 - removal of initial volumes
 - Siemens: 8 TRs
 - Philips: 8 TRs
 - GE DV25: 5 TRs
 - GE DV26: 16 TRs
 - normalization and demean
 - divide by the mean of each voxel, subtract 1, multiply by 100
 - regression
 - linear regression to remove quadratic trends and signals correlated with motion and mean time courses of cerebral white matter, ventricles, and whole brain, plus first derivatives (Power, et al., 2014; Satterthwaite, et al., 2012)
 - motion regression included 6 parameters plus derivatives and squares
 - frames with displacement > 0.3 mm were excluded from the regression (Power, et al., 2014)
 - temporal filtering
 - band-pass filtered between 0.009 and 0.08 Hz (Hallquist, et al., 2013)
 - pre-processed time courses were sampled onto the cortical surface
 - projecting 1mm into cortical gray matter along surface normal vector
 - motion censoring to reduce residual effects of head (Power, et al., 2012; Power, et al., 2014)
 - motion estimates filtered to attenuate signals (0.31 - 0.43 Hz) associated with respiration (18.6 - 25.7 respirations / minute)
 - time points with FD > 0.2 mm excluded from variance and correlation calculations
 - time periods with < 5 contiguous, sub-threshold time points also excluded
 - time points that were outliers in standard deviation across ROIs also excluded

- Regions of interest (ROI)
 - subcortical structures labeled with atlas-based segmentation (Fischl, et al., 2002)
 - cortical regions labeled with the Desikan atlas-based classification (Desikan, et al., 2006)
 - cortical regions labeled with the Destrieux atlas-based classification (Destrieux, et al., 2010)
 - functionally-defined parcels derived resting-state functional connectivity patterns (Gordon, et al., 2014)
- Functional connectivity analysis
 - seed-based, correlational approach (Van Dijk, et al., 2010)
 - adapted for cortical surface based analysis (Seibert and Brewer, 2011)
 - networks defined as pre-defined groups of cortical parcels (Gordon, et al., 2014)
 - e.g., default, fronto-parietal, dorsal attention, etc.
 - calculation of average within- and between-network mean correlation (Van Dijk, et al., 2010)
 - correlation between each pair of ROIs
 - Fisher transformed to z-statistics
 - averaged to provide measure of network correlation strength
 - calculation of correlation between each network and each subcortical ROI
- variance across time calculated for each ROI
 - magnitude of low frequency oscillations

References

- Andersson, J. L., Skare, S., & Ashburner, J. (2003). How to correct susceptibility distortions in spin-echo echo-planar images: application to diffusion tensor imaging. *NeuroImage*, 20(2), 870–888. [https://doi.org/10.1016/S1053-8119\(03\)00336-7](https://doi.org/10.1016/S1053-8119(03)00336-7).
- Cox, R.W. (1996) AFNI: software for analysis and visualization of functional magnetic resonance neuroimages. *Comput Biomed Res*, 29:162-73.
- Desikan, R.S., Segonne, F., Fischl, B., Quinn, B.T., Dickerson, B.C., Blacker, D., Buckner, R.L., Dale, A.M., Maguire, R.P., Hyman, B.T., Albert, M.S., Killiany, R.J. (2006) An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *Neuroimage*, 31:968-80.
- Destrieux C, Fischl B, Dale A, Halgren E. (2010) Automatic parcellation of human cortical gyri and sulci using standard anatomical nomenclature. *Neuroimage*. 2010 Oct 15;53(1):1-15.
- Fischl, B., Salat, D.H., Busa, E., Albert, M., Dieterich, M., Haselgrove, C., van der Kouwe, A., Killiany, R., Kennedy, D., Klaveness, S., Montillo, A., Makris, N., Rosen, B., Dale, A.M. (2002) Whole brain segmentation: automated labeling of neuroanatomical structures in the human brain. *Neuron*, 33:341-55.
- Gordon, E.M., Laumann, T.O., Adeyemo, B., Huckins, J.F., Kelley, W.M., Petersen, S.E. (2014) Generation and Evaluation of a Cortical Area Parcellation from Resting-State Correlations. *Cereb Cortex*.
- Hallquist, M.N., Hwang, K., Luna, B. (2013) The nuisance of nuisance regression: spectral misspecification in a common approach to resting-state fMRI preprocessing reintroduces noise and obscures functional connectivity. *Neuroimage*, 82:208-25.
- Holland, D., Kuperman, J.M., Dale, A.M. (2010) Efficient correction of inhomogeneous static magnetic field-induced distortion in Echo Planar Imaging. *Neuroimage*, 50:175-83.
- Jovicich, J., Czanner, S., Greve, D., Haley, E., van der Kouwe, A., Gollub, R., Kennedy, D., Schmitt, F., Brown, G., Macfall, J., Fischl, B., Dale, A. (2006) Reliability in multi-site structural MRI studies: effects of gradient non-linearity correction on phantom and human data. *Neuroimage*, 30:436-43.
- Power, J.D., Barnes, K.A., Snyder, A.Z., Schlaggar, B.L., Petersen, S.E. (2012) Spurious but systematic correlations in functional connectivity MRI networks arise from subject motion. *Neuroimage*, 59:2142-54.
- Power, J.D., Mitra, A., Laumann, T.O., Snyder, A.Z., Schlaggar, B.L., Petersen, S.E. (2014) Methods to detect, characterize, and remove motion artifact in resting state fMRI. *Neuroimage*, 84:320-41.
- Satterthwaite, T.D., Wolf, D.H., Loughhead, J., Ruparel, K., Elliott, M.A., Hakonarson, H., Gur, R.C., Gur, R.E. (2012) Impact of in-scanner head motion on multiple measures of functional connectivity: relevance for studies of neurodevelopment in youth. *Neuroimage*, 60:623-32.
- Seibert, T.M., Brewer, J.B. (2011) Default network correlations analyzed on native surfaces. *J Neurosci Methods*, 198:301-11.

Smith, S. M., Jenkinson, M., Woolrich, M. W., Beckmann, C. F., Behrens, T. E., Johansen-Berg, H., Bannister, P. R., De Luca, M., Drobnjak, I., Flitney, D. E., Niazy, R. K., Saunders, J., Vickers, J., Zhang, Y., De Stefano, N., Brady, J. M., & Matthews, P. M. (2004). Advances in functional and structural MR image analysis and implementation as FSL. *NeuroImage*, 23 Suppl 1, S208–S219. <https://doi.org/10.1016/j.neuroimage.2004.07.051>.

Van Dijk, K.R., Hedden, T., Venkataraman, A., Evans, K.C., Lazar, S.W., Buckner, R.L. (2010) Intrinsic functional connectivity as a tool for human connectomics: theory, properties, and optimization. *J Neurophysiol*, 103:297-321.

Wells, W.M., 3rd, Viola, P., Atsumi, H., Nakajima, S., Kikinis, R. (1996) Multi-modal volume registration by maximization of mutual information. *Med Image Anal*, 1:35-51.

ABCD Study[®], Teen Brains. Today's Science. Brighter Future[®]. and the ABCD Study Logo are registered marks of the U.S. Department of Health & Human Services (HHS). Adolescent Brain Cognitive DevelopmentSM Study is a service mark of the U.S. Department of Health & Human Services (HHS).